

# Benchmark No. 11

## Management of Behaviour and Cognitive Impairment



**British Association of  
Neuroscience Nurses**



# Neuroscience Safe Staffing Benchmark Statements

Copyright © 2020 British Association of Neuroscience Nurses. All rights reserved.

This PDF 1<sup>st</sup> edition printed 2020 in the United Kingdom (available online).

A catalogue record for this book is available from the British Library.

ISBN 978-1-911059-18-9

No part of this book shall be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or by any information retrieval system without written permission of the publisher.

Published by the British Association of Neuroscience Nurses

For more copies of this book, please email: [info@bann.org.uk](mailto:info@bann.org.uk)

Designed and Set by the British Association of Neuroscience Nurses

[www.bann.org.uk](http://www.bann.org.uk)

Printed in the United Kingdom

***Although every precaution has been taken in the preparation of this publication, the publisher and authors assume no responsibility for errors or omissions. Neither is any liability assumed for damages resulting from the use of this information contained***



## History

The Neuroscience Nursing Benchmarking Group (NNBG) was established in the 1990`s as a result of increasing concerns over inconsistencies in practices as part of a subsidiary of BANN. The group aims to improve on the quality of care by comparing and sharing practice with each other, and set explicit standards for comparison of current practice against the ideal standard. The group is committed to searching for the best evidence related to specific areas of neuroscience practice. Membership of the group consists of representatives from neuroscience units within the UK and Ireland, together with educational colleagues from both the NHS/HSC and Higher Educational Institutes. The group is further subdivided into regions and the first edition of this benchmark was developed by the North West regional group of the NNBG in 2006.

In 2016, the NNBG consolidated back into BANN and further information about NNBG can be found on the BANN website [www.BANN.org.uk](http://www.BANN.org.uk) . This second edition of the benchmark has been developed by the restructured NNBG working group under BANN.

BANN would like to acknowledge the leadership and significant contribution made by the NNBG, and all its contributors, to neuroscience nursing over the years.

## Benchmark No.11 Management of Behaviour and Cognitive Impairment

### KEY POINTS

- Following a full risk assessment an individualised multi-disciplinary care plan must be implemented and evaluated specific to all aspects of care relating to the use of any restrictive interventions.
- If the use of restrictive interventions amounts to a deprivation of liberty, then the relevant referrals must be made under the Deprivation of Liberty Safeguards (MCA, 2005, Mental Capacity (Amendment) Act 2019).
- Accurate documentation includes the clinical need for the physical intervention, the type of intervention employed, the date and time that the intervention was implemented, reviewed and discontinued.
- The least restrictive therapy should be chosen for the shortest period of time.
- The patient must be reassessed at regular intervals or when their health needs change.
- Patients and relatives are included in the decision-making process and on-going management wherever possible. All verbal and written information must be current; evidence based and is documented in the patient's notes.
- Pharmacological interventions are only considered when all other factors have been considered in the management of behaviour and cognitive impairment.
- Documentation must include the drug, the dose the mode of delivery, time frame and potential contra-indications.
- Clinical staff are provided with a structured competency-based training and education programme for the care of the patient.
- The care plan is evidence based, dated and reviewed within the last two years and updated accordingly.
- Best interest meetings with family/carers are clearly documented.

FACTOR 1 – Documentation

STATEMENT OF BEST PRACTICE	EVIDENCE & REFERENCES	ACHIEVED	NOT ACHIEVED	VARIABLES
<p>1.0 A full risk assessment of the patient is performed and documented prior to the use of any restrictive interventions. This must include a comprehensive assessment addressing:</p> <ul style="list-style-type: none"> <li>a. The cause and the triggers for the altered behaviour i.e. physiological (dementia, learning disabilities, possible drug interactions), psychiatric (including substance abuse and drug history) or neurological (brain injury, infection).</li> <li>b. Complete history of pre-morbid drug history including non-prescribed drugs, alcohol and substance, benzodiazepine use.</li> <li>c. Assessment of mental capacity and the ability to consent to treatment.</li> <li>d. The use of a validated assessment scale to establish a baseline level of agitation or aggressive behaviour i.e. Mini-Mental State Examination (MMSE), Modified Overt Aggression Scale (MOAS). Richmond Agitation Sedation scale (RAS).</li> <li>e. The diagnosis of possible delirium is made using a recognised assessment tool.</li> <li>f. There is evidence of the use of alternative approaches to the management of the behaviour i.e. environmental modifications, behavioural or exercise programmes.</li> <li>g. Pharmacological interventions and escalation guidelines are available.</li> <li>h. There is an identified schedule for reassessment of the intervention.</li> </ul>	<p>NPSA 2015</p> <p>DoH, 2005</p> <p>Braine, 2005</p> <p>Mattes, 2010.</p> <p>Rasheed <i>et al</i> 2019</p>			

STATEMENT OF BEST PRACTICE	EVIDENCE & REFERENCES	ACHIEVED	NOT ACHIEVED	VARIABLES
1.1 Following a multi-disciplinary team assessment an individualised care plan is implemented and regularly re-evaluated in accordance with the patients changing healthcare needs.				
1.2 An inter-disciplinary care plan includes the following issues: <ul style="list-style-type: none"> <li>a) Consideration of pre-disposing or trigger factors that may be contributing to the behaviours. e.g. Environment, time of day, pain, hunger, etc.</li> <li>b) Dialogue with the family/carers on the patient's need for restrictive interventions.</li> <li>c) Daily multi-disciplinary re-evaluations of care delivered, including consultation with a pharmacist.</li> <li>d) Recognition that the least restrictive intervention has been used for the shortest period of time.</li> <li>e) The date and time that the intervention was implemented, reviewed and discontinued.</li> <li>f) Mental Capacity assessment (if appropriate).</li> <li>g) Deprivation of Liberty documentation (if appropriate).</li> </ul>	RCN, 2017         Ridley & Leitch, 2019			
1.3 An escalation plan is in place for continuing management.				

**FACTOR 2 – Protocol**

STATEMENT OF BEST PRACTICE	EVIDENCE & REFERENCES	ACHIEVED	NOT ACHIEVED	VARIABLES
<p>2.0 Evidence based guidelines/policies are available for the management of behaviour and cognitive impairment. For patients who lack capacity this will include:</p> <ul style="list-style-type: none"> <li>a) The most appropriate intervention for managing the patient’s behaviour.</li> <li>b) Clinical holding policy.</li> <li>c) Guidelines for debriefing of staff post-incident.</li> <li>d) Family/carers have the opportunity to be included in the decision-making process and on-going management wherever possible</li> </ul>	<p>NPSA 2015</p>			
<p>2.1 <b>Pharmacological Interventions</b></p> <p>Protocol and guidelines Staff providing direct patient care have knowledge of the following:</p> <ul style="list-style-type: none"> <li>a) Administration of covert medications policy.</li> <li>b) Rapid Tranquilisation</li> <li>c) The patient is prescribed and administered the appropriate medication to meet their needs.</li> <li>d) The effectiveness and response to the pharmacological intervention is monitored.</li> <li>e) Contraindication with other medications is documented.</li> <li>f) The potential risks, complications and side effects of prescribed medications is identified e.g., extra pyramidal symptoms, dystonia, anti-cholinergic effects.</li> </ul>	<p>Bourne, 2008 Chew, 2009  Ridley &amp; Leitch, 2019</p>			

STATEMENT OF BEST PRACTICE	EVIDENCE & REFERENCES	ACHIEVED	NOT ACHIEVED	VARIABLES
<p><b>2.2 Physical Interventions</b></p> <p>Staff providing direct patient care have knowledge of the following:</p> <ul style="list-style-type: none"> <li>a) Conflict resolution.</li> <li>b) De-escalation skills.</li> <li>c) Personal risks associated with clinical holding.</li> <li>d) Identification of potential risks to the patient of clinical holding.</li> <li>e) Level of clinical holding is proportionate to the patient's behaviour.</li> <li>f) Documentation of tissue integrity and patient hygiene.</li> <li>g) Falls Risk Assessment</li> </ul>	<p>MCA, 2005.</p> <p>MCA, Liberty Protection safeguards 2020</p> <p>DOL's, 2015</p> <p>Cleary, 2015</p> <p>NPSA. (2015).</p>			



FACTOR 3 – Education

STATEMENT OF BEST PRACTICE	EVIDENCE & REFERENCES	ACHIEVED	NOT ACHIEVED	VARIABLES
<p>3.0 All healthcare professionals applying restrictive interventions are provided with local training.</p> <p>Training includes:</p> <ul style="list-style-type: none"> <li>a) Assessment and recognition of behaviours</li> <li>b) Conflict Resolution training.</li> <li>c) Breakaway techniques.</li> <li>d) Importance of de-escalation, distraction, interaction, verbal and non-verbal communication.</li> <li>e) Completion of a comprehensive risk assessment.</li> <li>f) Defining what is classified as ‘Physical Restraint’</li> <li>g) Defining what is classified as ‘Clinical holding’.</li> <li>h) Awareness of personal professional responsibilities and legal accountability when employing restrictive interventions.</li> <li>i) Recognition of the potential risks, personal safety and complications associated with restrictive interventions</li> <li>j) Consideration of the legal and ethical issues underpinning enhanced care.</li> <li>k) Importance of limiting sensory under-load or over-load.</li> <li>l) Identification of techniques and equipment to maintain patient safety, for example: <ul style="list-style-type: none"> <li>• Mechanical restraints – mittens, splints, bedrails, tilt back chairs, seat belts, lap straps intended to provide postural support.</li> <li>• Electronic Surveillance – tagging, pressure pads, door alarms.</li> </ul> </li> </ul>	<p>DoH, 2005, 2012. 2014 2015</p> <p>NPSA. 2015.</p> <p>Cleary, 2015</p> <p>NMC, 2015</p> <p>Restraint reduction network, 2019</p> <p>Vindola-Padros <i>et al</i> 2018</p>			

STATEMENT OF BEST PRACTICE	EVIDENCE & REFERENCES	ACHIEVED	NOT ACHIEVED	VARIABLES
<ul style="list-style-type: none"> <li>• Psychological restraint – repeated verbal commands.</li> <li>• One to One enhanced care.</li> <li>• Seclusion – confinement and isolation of a patient away from the other patients.</li> </ul> <p>m) Nursing care relevant to the restrictive intervention e.g., tissue viability.</p> <p>n) Nonpharmacological sleep promotion strategies.</p>				
<p>3.1 Staff have knowledge of delirium assessment using a recognised tool i.e. Confusion Assessment Method (CAM), confusion assessment method for the intensive care unit (CAM-ICU), Intensive care delirium screening checklist (ICDSC).</p>	<p>Bourne, 2008</p>			
<p>3.2 Staff can demonstrate awareness of the legal and ethical implications of using restrictive interventions associated with mental health and mental capacity:</p> <ul style="list-style-type: none"> <li>a) Mental Health Act (1987, 2007) Great Britain, Mental Health Act (2001) Ireland</li> <li>b) Adults with Incapacity Scotland (2000) Mental Capacity Act England, Wales, Ireland &amp; Northern Ireland (DH, 2005)</li> <li>c) Human Rights Act (1998)</li> <li>d) DoL's (2015) &amp; Vulnerable Adult Order</li> <li>e) 'Best interest' decision making</li> </ul>				

**FACTOR 4 – Patient Information**

STATEMENT OF BEST PRACTICE	EVIDENCE & REFERENCES	ACHIEVED	NOT ACHIEVED	VARIABLES
4.0 Patient information is available and reviewed in accordance with local policy.				
4.1 Patients/carers must be given current evidence based verbal and written information including: <ul style="list-style-type: none"> <li>a. Rationale for the intervention (physical, pharmacological).</li> <li>b. How often the patient will be reviewed</li> <li>c. Possible complications</li> <li>d. Risks and benefits</li> </ul>	DH, 2005, 2012, 2014.			
4.2 Any information verbal /written that is given to the patient/carers is documented in the patients notes				

## References

- Braine, M.E. (2005). The management of challenging behaviour and cognitive impairment. *British Journal of Neuroscience Nursing* 1(2): 67–74
- Bourne RS. (2008). Drug treatment of delirium: Past, present and future. *Journal of Psychosomatic Research* ; 65 (3): 273 – 282. [https://www.jpsychores.com/article/S0022-3999\(08\)00246-8/fulltext](https://www.jpsychores.com/article/S0022-3999(08)00246-8/fulltext)
- Chew E, Zafonte RD: Pharmacological management of neurobehavioral disorders following traumatic brain injury—a state-of-the-art review. *J Rehabil Res Dev* 2009, 46:851-879.
- Cleary KK, Prescott K. (2015). The use of Physical Restraints in Acute and Long-term care: n update Review of the evidence, regulations , ethics and legality. *The journal of acute care, physical therapy.* 6(1): 8-15.
- Department of Health (2005) Deprivation of Liberty Safeguards policy document, Department of Health. [DoLS guidelines](#)
- Department of Health (2015) *Deprivation of liberty safeguards – information and resources*, London, DH. Available from: [www.gov.uk/government/publications/deprivation-of-liberty-safeguards-forms-and-guidance](http://www.gov.uk/government/publications/deprivation-of-liberty-safeguards-forms-and-guidance)
- Department of Health (2012) *Transforming Care: a national response to Winterbourne View Hospital*, London: DH.
- Department of Health (2014) *Positive and Proactive Care: reducing the need for restrictive interventions*, London: DH. Department of Health, Skills for care and skills for health. *A positive and proactive workforce*, London: DH. [www.skillsforcare.org.uk/Topics/Restrictivepractices/Restrictive-practices.aspx](http://www.skillsforcare.org.uk/Topics/Restrictivepractices/Restrictive-practices.aspx)
- Department of Health, Social Services and Public Safety Northern Ireland (2003) *Reference Guide to Consent for Examination, Treatment or Care*, Belfast, DHSSPS.
- Department of Health, Social Services and Public Safety Northern Ireland (2010) Circular HSC/ MHDP – MHU 1 /10 – revised. *Deprivation of Liberty Safeguards. (DOLS) – Interim Guidance*, Available from: [www.dhsspsni.gov.uk/publications/deprivation-liberty-safeguardsdols-%E2%80%93-interim-guidance](http://www.dhsspsni.gov.uk/publications/deprivation-liberty-safeguardsdols-%E2%80%93-interim-guidance).
- Mattes, J. A. (2010). Suggested Improvements to the Overt Aggression Scale-Modified. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 22(1), 123-123.
- Mental Capacity Act 2005 and associated Code of Practice: London, HMSO. <http://www.justice.gov.uk/guidance/protecting-the-vulnerable/mentalcapacity-act/index.htm> 14. [Mental Health Act Policy.](#)
- Mental Capacity Act 2005, Deprivation of Liberty Safeguards code of Practice (To be superseded by Liberty Protection safeguards 2020)
- Mental Health Act (2015). <https://www.legislation.gov.uk/ukpga/2005/9>. The Stationery Office London.
- National Institute for Health and Care Excellence NICE (2015)., Violence and aggression: short term management in mental health, health and community settings. Clinical Guideline CG103. <https://www.nice.org.uk/guidance/cg103>
- National Institute for Health and Care Excellence (2015) *Challenging Behaviour and Learning Disabilities: Prevention and Interventions for People with Learning Disabilities whose behaviour Challenges* (NG11). Available from: [www.nice.org.uk](http://www.nice.org.uk)
- NPSA. (2015). Introduction to Challenging Behaviour - Meeting needs and reducing distress. [www.nhs.protect.nhs.uk/reducingdistress](http://www.nhs.protect.nhs.uk/reducingdistress)

Northern Ireland Assembly (2016) *Mental Capacity Act (Northern Ireland)*.

[www.legislation.gov.uk/nia/2016/18/contents/enacted](http://www.legislation.gov.uk/nia/2016/18/contents/enacted)

Ridley J, Leitch S (2019) Restraint Reduction Network, RRN. Ethical training standards to protect human rights and minimise restrictive practices. British Institute of learning disabilities.


Rasheed, AM. Amirah MF. Parameaswari, Marwan I, Alharthy, Abdulrhman Mohammad F. Ramsay (2019). Sedation Scale and Richmond Agitation Sedation Scale: A cross sectional Study. Dimensions of Critical care Nursing. Vol 38. 2 p90-99

Royal College of Nursing (2017). Three steps to Positive Practice. A rights based approach when considering and reviewing the use of restrictive interventions.

Scottish Executive Health Department (2001) *Adults with Incapacity (Scotland) Act 2000* ISBN 0 10 590005 2. HMSO. Edinburgh, Scottish Government.

Scottish Government (2003). *Mental Health, (Care and Treatment) (Scotland) Act*, Edinburgh, Scottish Government.

Vindrola-Padros, C, Swart N, McIntosh M, Crowe S, Morris S, Fulop N. (2018). One to One specialing and sitters in acute care hospital: A scoping review. International Journal of Nursing Studies. 84, 61-77



**Benchmark No. 4 (2<sup>nd</sup> Ed)**  
**Management of Behaviour and Cognitive Impairment**

ISBN 9781911059189



9 781911 059189